



Revisiting Stress “Deafness” in European Portuguese – An ERP study

Shuang Lu, Susana Correia, Rita Jerónimo,
Marina Vigário, Sónia Frota

Speech Prosody 2016,
Boston University

Introduction

- Lexical stress refers to the prominent syllable in a word
 - Fixed stress: stress always falls on a particular position
 - E.g., Finnish, Polish, Turkish
 - Variable stress: the position of stress in a word is not predictable
 - E.g., English, Spanish, German
 - Processing of word stress is particularly relevant in such languages
 - Minimal pairs that only differ in stress pattern (e.g., insight /^linsart/ vs. incite/ in^lsart/ in English)
- Speakers of languages with variable stress are **better** than speakers of languages with fixed stress in distinguishing non-words that differ only in stress pattern (e.g., Domahs et al., 2012; Dupoux et al., 1997, 2001, 2008; Peperkamp et al., 2010; Rahmani et al. 2015)

Introduction

- Lexical stress is signaled by phonetic cues:
 - E.g., Duration, F0, intensity, vowel quality
- Languages differ in the weighing of phonetic cues
 - **English**: F0 contour > intensity, duration, vowel quality (Beckman, 1986; Fry, 1958)
 - **Catalan**: duration, spectral balance, vowel quality (Astruc & Prieto, 2006)
- The absence of certain cues may influence listeners' perception of stress

Introduction

◦ European Portuguese (EP): Variable stress

◦ Vowel reduction: primary cue

◦ Behavioral studies (ABX, Sequence Recall Task): without vowel reduction cues → **stress “deafness” effect** similar to that found for languages with fixed stress (Correia et al., 2015)

◦ Duration: main prosodic cue (Delgado-Martins, 1977; Andrade & Viana, 1989) in the absence of vowel reduction

◦ But not sufficient for the processing of stress contrasts

◦ Pitch: low correlate of stress, given sparse distribution of pitch accents (Frota, 2014)

◦ Frequency asymmetry: Trochee > iamb (Types - Vigário et al. 2010)

◦ No previous study has been conducted to examine the **unintentional processing of stress** by native EP speakers

The Current study

Research questions:


- Can native speakers of EP unintentionally discriminate stress in the absence of vowel reduction?
 - Previous research suggested that discrimination may occur at the unintentional level, but not at the intentional/behavioral level (Tremblay et al., 1998)
- Will native speakers of EP show asymmetric effects to the two stress patterns in EP?
 - Frequency based on type – advantage for trochee

The Current study

- ERP experiment: passive oddball paradigm

- Two ERP components:

- Mismatch Negativity (MMN): negative wave elicited by auditory stimuli that are infrequently presented and deviate from a frequently presented standard stimuli : X X X X Y X X X (Näätänen et al., 2004)



- Late negativity: negative wave that occurs around 350-600ms after the onset of deviant stimuli; has been associated with neural processes of auditory rule extraction (Zachau et al. 2005)

Methods

Participants

- 24 native speakers of EP (6 males, 18 females)
- Between the ages of 18-32 ($M=21.92$, $SD=3.97$)
- Right-handed according to the Edinburgh Handedness inventory
- Normal vision and hearing
- No history of speech or neurological impairment

Methods

Stimuli

- Disyllables [bubu] with either a trochaic or an iambic stress in the absence of vowel reduction
- Each of the stress patterns was produced twice by a female native speaker of EP [l'bubu]1, [l'bubu]2, [bu'bu]1, and [bu'bu]2
- Mean durations: trochaic tokens – 872ms
iambic tokens – 873ms
- The first 100 millisecond of [l'bubu]1, [l'bubu]2, and [bu'bu]2 were replaced by the first 100 millisecond of [bu'bu]1, to control the acoustic onset differences.
- After the manipulation no pitch discontinuity was observed
- Three native EP speakers who did not participate in the ERP experiment judged all the stimuli as perceptually natural.





Figure 1: Spectrograms of the trochaic and iambic stress patterns

Methods

o Procedure

- o Iambic block: trochaic standards (250 times) vs. iambic deviants (50 times) 
- o Trochaic block: iambic standards (250 times) vs. trochaic deviants (50 times)
- o 600 trials (250 x 2 tokens + 50 x 2 tokens)
- o Pseudo-random order, with at least 2 standards preceding each deviant
- o Offset-to-onset inter-stimulus interval randomly varied between 800~850ms

Methods

○ Procedure

- Participants were watching a muted movie (*The Gold Rush* by Charlie Chaplin) in a sound-attenuating booth, while the stimuli were presented through a loudspeaker
- Participants were asked to ignore the sounds and focus on the movie.
 - They received comprehension questions regarding the movie after each block
- E-Prime 2.0 software was used
- 29 Ag/AgCl scalp electrodes were recorded (Easy-Cap, SynAmps1, NeuroScan)
- After EEG data processing, on average 96 trials for each stimulus type were included in data analysis

10


Results

- Both the Trochaic and Iambic conditions yielded MMN and late negativity


a. Trochee	300-400	400-500	500-600	600-700	700-800	800-900
Disc	***			*		
Ante	***	***	**	**	***	***
Hemi		*	**	*	**	**
Disc × Hemi		*				*
Disc × Ante				*		
Hemi × Group		**	*			
Disc × Hemi × Ante	*	**	**	**	*	
Disc × Hemi × Ante × Group			**	*		
b. Iamb	300-400	400-500	500-600	600-700	700-800	800-900
Disc	***	***	***	***	**	*
Ante	***	***	***	***	***	***
Hemi		*				
Group						*
Disc × Ante					***	**
Hemi × Ante			*	*		*
Hemi × Ante × Group			*	**	**	**

Figure 2: Grand averages

a. Trochee



b. Iamb



Results

- The MMN and late negativity components in the iambic condition span over a larger temporal window than in the trochaic condition:




Figure 3: *Grand-average difference waves*

Follow-up: behavioral study

- o ABX discrimination task (as in Correia et al., 2015): Goal – replicate previous findings of a **stress “deafness” effect**

- o Participants

- o 21 native speakers of EP who had participated in the ERP experiment

- o Stimuli

- o 9 pairs of disyllabic nonsense words with trochaic and iambic stress, *without vowel reduction*, e.g., [l̥midu], [mi̯du]

- o Procedure

- o A,B were always produced by two female speakers (same or different) and X by a male speaker

- o AB/BA counterbalanced

- o The participant’s task is to respond whether X=A or X=B

13

Follow-up: behavioral study

Results

- Mean error rate: 21.4%

- A stress “deafness” effect similar to French (Dupoux et al., 1997)

Stress ABX	French	Spanish	EP
Error rate	19%	4%	21%

- Replicates the findings from Correia et al. (2015):


- Same task, different participants

- EP – 21% error rate (vs. 5% with a phoneme contrast)

Follow-up: behavioral study

Results

- A difference was found between lamb and Trochee
- Lower error rate when $X=\text{lamb}$, both when X is the same or different from the preceding stimuli



- RTs faster when $X=\text{lamb}$

Discussion

- o Can native speakers of EP unintentionally discriminate stress in the absence of vowel reduction?
 - o **YES!** Trochaic and iambic conditions yielded MMN and late negativity
 - o However, previous behavioral studies demonstrated **a stress “deafness” effect**; ABX results replicated in the current study
 - o Suggests listeners are using some **acoustics-based strategy**, relying on phonetics similarity/difference in the ERP task
 - o Results from the ABX tasks (more variation in the stimuli) show that the **acoustic cues are not that robust** in EP (or French, unlike Spanish) > EP listeners have more difficulty, but do not fail
 - o Results from the Sequence Recall Task tap into a **more abstract/phonological representation** (Dupoux et al. 2001; Rahmani et al. 2015) > EP listeners fail (in the presence of prosodic cues only)

Discussion

- Perceptual discrimination may occur at the unintentional level, but not at the intentional/behavioral level
- Consistent with findings for Polish: Polish speakers are “deaf” to stress manipulations in a behavioral task but show different neural responses towards default stress vs. exceptions (Domahs et al. 2012)
- Polish is a fixed stress language (penult stress), with well-defined exceptions



- Neural response to stress contrasts even in languages with (some degree of) stress “deafness”

Discussion

◦ Will native speakers of EP show asymmetric effects to the two stress patterns?

◦ **YES!** They are more sensitive to the **iambic** pattern

◦ Arguably contrary to the frequency distribution of stress patterns in EP (type) and previous literature on other languages using ERP

◦ E.g. native speakers of Russian are more sensitive to the trochaic stress pattern, which faithfully represents the frequency asymmetries of the stress patterns in the language (Molczanow et al. 2013).

◦ But frequency distribution changes if based on token, if monosyllabic words are included with final stress words (Vigário et al. 2010)

◦ Some ERP studies show less sensitivity to the most frequent (or default) pattern in the language (Friederici et al. 2007)

Discussion

- Will native speakers of EP show asymmetric effects to the two stress patterns?
 - **YES!** They are more sensitive to the **iambic** pattern
 - Consistent with behavioral results in adult and infant studies
 - Adult studies show lower error rates when X=lamb in the ABX task
 - A study on infants' perception of stress showed that 5-6 month old EP-learning infants prefer the iambic to the trochaic pattern (Butler et al. 2015)
- The current findings
 - Differences in how stress information is encoded in a perception task
 - EP speakers are more sensitive to iambic stress



Thank you !
Obrigada!
谢谢 !

Contacts



Shuang Lu

shuanglu2013@outlook.com

School of Foreign Languages,
Renmin University of China, China



Susana Correia

correia.smd@gmail.com

Centre of Linguistics,
Universidade de Lisboa, Portugal



Rita Jerónimo

rita.jeronimo@iscte.pt

Instituto Universitário de Lisboa
(ISCTE-IUL), CIS-IUL, Lisboa, Portugal

Marina Vigário

marina.vigario@mail.telepac.pt

Centre of Linguistics,
Universidade de Lisboa,
Portugal



Sónia Frota

sonia.frota@mail.telepac.pt

Centre of Linguistics,
Universidade de Lisboa,
Portugal

References

- o A., Friederici, M. Friedrich, and A. Christophe, “Brain Responses in 4 Month-Old Infants Are Already Language Specific”, *Current Biology*, 17(14), pp. 1208-1211, 2007.
- o D. B. Fry, “Experiments in the Perception of Stress,” *Language and Speech*, 1, pp. 126-152, 1958.
- o E. Andrade, and M. C. Viana, “Ainda sobre o ritmo e o acento em português,” *Act. do IX Enc. Nac. da APL*, pp. 3-15, 1989.
- o E. Dupoux, C. Pallier, N. Sebastian, and J. Mehler, “A Destressing 'Deafness' in French?” *Journal of Memory and Language*, 36, pp. 406-421, 1997.
- o E. Dupoux, N. Sebastián-Gallés, E. Navarrete, and S. Peperkamp, “Persistent Stress ‘Deafness’: The Case of French Learners of Spanish,” *Cognition*, 106, pp. 682–706, 2008.
- o E. Dupoux, S. Peperkamp, and N. Sebastián-Gallés, “A Robust Method to Study Stress ‘deafness’,” *Journal of Acoustical Society of America*, 110, pp. 1606–1618, 2001.
- o H. Rahmani, T. Rietveld, and C. Gussenhoven, “Stress “Deafness” Reveals Absence of Lexical Marking of Stress or Tone in the Adult Grammar”, *PLOS One*, 2015.
- o J. Bulter, S. Cortês, S. Correia, E. Uysal, M. Vigário, and S. Frota, “Early perception of lexical stress by European Portuguese-learning infants,” *Talk presented at PaPE2015 - Phonetics and Phonology in Europe*, Cambridge: University of Cambridge, 2015.
- o J. Molczanow, U. Domahs, J. Knaus, and R. Wiese, “The Lexical Representation of Word Stress in Russian: Evidence from Event-related Potentials,” *The Mental Lexicon*, 8, pp. 164–194, 2013.

References

- K. Tremblay, N. Kraus, and T. McGee, "The Time Course of Auditory Perceptual Learning: Neurophysiological Changes during Speech-Sound Training," *Neuroreport*, 9(16), pp. 3557-3560, 1998.
- L. Astruc and P. Prieto, "Stress and Accent: Acoustic correlates of Metrical Prominence in Catalan," In: *ITRW on Experimental Linguistics*, pp. 73-76, 2006.
- M. E. Beckman, *Stress and Non-stress Accent*. Dordrecht, the Netherlands: Foris Publications, 1986.
- M. R. Delgado-Martins, *Aspects de l'accent en portugais. Voyelles toniques et atones*. PhD Dissertation of 3rd cycle. University of Strasbourg, 1977.
- M. Vigário, S. Frota, and F. Martins. "A Frequência que Conta na Aquisição da Fonologia." *Textos Seleccionados. XXV Encontro Nacional da APL*. 2010
- R. Näätänen, S. Pakarinen, T. Rinne, and R. Takegata, "The Mismatch Negativity (MMN): Towards the Optimal Paradigm," *Clinical Neurophysiology*, 115, pp. 140–144, 2004.
- S. Correia, J. Butler, M. Vigário and S. Frota, "A Stress "Deafness" Effect in European Portuguese," *Language and Speech*, 58(1), pp. 24-47, 2015.
- S. Frota, "The Intonational Phonology of European Portuguese". In Sun-Ah Jun (Ed.). *Prosodic Typology II*. Oxford: Oxford University Press. Pp 6-42, 2014.
- S. Peperkamp, I. Vendelin, and E. Dupoux, "Perception of Predictable Stress: A Cross-linguistic Investigation," *Journal of Phonetics*, 38, pp. 422–430, 2010.
- S. Zachau, T. Rinker, B. Körner, G. Kohls, V. Maas, K. Henninghausen, and M. Schecker, "Extracting Rules: Early and Late Mismatch Negativity to Tone Patterns," *Neuroreport*, 16, pp. 2015-2019, 2005.
- U. Domahs, J. Knaus, P. Orzechowska, and R. Wiese, "Stress "Deafness" in a Language with Fixed Word Stress: an ERP Study on Polish," *Frontiers in Psychology*, 3:439, 2012.